CLEAN COPY OF THE AMENDED CLAIMS

4. (Thrice Amended) A monoclonal antibody or recombinant antibody fragment having the capability of binding a human gamma-carboxylated osteocalcin fragment, comprising the amino acid sequence

in which at least one of the gluramic acids in positions 17, 21 and 24 is gamma-carboxylated, characterized by the specificity to epitopes that have been identified on the gamma-carboxylated fragment of osteocalcin, wherein said fragment spans either

- i) from the amino acid in position 7 to the amino acid in position 30, or
- ii) from the amino acid in position 6 to the amino acid in position 30 of the amino acid sequence of SEQ ID NO:2, and that all three glutamic acids in the positions 17, 21 and 24 of said sequence are gamma-carboxylated.
- 6. (Amended) A non-competitive immunoassay for quantitative determination of a gamma-carboxylated osteocalcin fragment, comprising the amino acid sequence

Serial No.: 09/462,931 19 April 2001 Page 3

in which at least one of the glutamic acids in positions 17, 21 and 24 is gamma-carboxylated, characterized in that a sample containing said fragment is exposed to two monoclonal antibodies or recombinant antibody fragments which bind said gamma-carboxylatedosteocalcin fragment, said monoclonal antibodies or recombinant antibody fragments are specific to epitopes that have been identified on the gamma-carboxylated fragment of osteocalcin, wherein said fragment spans either

i) from the amino acid in poisition 7 to the amino acid in position 30, or

ii) from the amino acid in position 6 to the mino acid in position 30 of the amino acid sequence of SEQ ID NO:2, and that all three glutamic acids in the positions 17, 21 and 24 of said sequence are gamma-carboxylated.

